

Appl. No. : 10/063,540  
Filed : May 2, 2002

### REMARKS

Claim 1 has been amended to add the term "isolated." Support for this amendment can be found at, for example, page 47, lines 26-35. Claims 1-5 remain present for examination. Applicants thank the Examiner for the review of the instant application and withdrawing the objection to the title and specification, acceptance of the copy of the sequence listing, Applicants' request for correction of inventorship, and the submission of the IDS dated September 29, 2004. Applicants acknowledge the Examiner's withdrawal of the rejection of Claims 1-6 under 35 U.S.C. § 112, second paragraph, as being indefinite, and the Claims 1-5 under 35 U.S.C. § 103(a). The rejections of the presently pending claims are respectfully traversed.

#### **Priority**

The PTO asserts that because the disclosure of PCT/US00/23328 is not enabling for the instant invention, the filing date of the present application is May 2, 2002.

Applicants have previously listed the priority information for the instant application in a Preliminary Amendment mailed September 3, 2002. Applicants submit that for the reasons stated herein, the claimed antibodies are enabled by the disclosure of PCT/US00/23328 and have a credible, substantial, and specific utility. Applicants maintain that the present application is entitled to at least the priority date of August 24, 2000.

#### **Finality of the Rejection is Premature**

The PTO maintains its rejection of Claims 1-5 under 35 U.S.C. § 101 as lacking utility, stating that the reasons are set forth in the previous Office Action at pages 3-8. However, instead of relying on the previously cited references and arguments which are not relevant to the instant application, the PTO has adopted new arguments, and has cited three new references for support. Because the PTO has changed its arguments, and is no longer relying on the references cited in the previous Office Action, Applicants submit that the finality of the present rejection is improper. M.P.E.P. § 706.07 makes this point clear:

Before final rejection is in order a clear issue should be developed between the examiner and applicant. ... Switching from one subject matter to another in the claims presented by applicant in successive amendments, or from one set of references to another by the examiner in rejecting in successive actions claims of substantially the same subject matter, will alike tend to defeat attaining the goal of

Appl. No. : 10/063,540  
Filed : May 2, 2002

reaching a clearly defined issue for an early termination, i.e., either an allowance of the application or a final rejection. ... The applicant ... should receive the cooperation of the examiner ... and not be prematurely cut off in the prosecution of his or her application. ... The examiner should never lose sight of the fact that in every case the applicant is entitled to a full and fair hearing, and that a clear issue between applicant and examiner should be developed, if possible, before appeal.

Because the PTO is relying on new references and new arguments, Applicants should be given the opportunity to submit references and arguments in response. Finality is improper because a clear issue has not been developed for Appeal until Applicants have had an opportunity to place their arguments and supporting documents in the record. Applicants therefore request that the finality of the rejection be withdrawn, or at a minimum, that the arguments and references cited herein be entered into the record for purposes of appeal.

#### **Rejection Under 35 U.S.C. §101 – Utility**

The PTO maintains its rejection of Claims 1-5 under 35 U.S.C. § 101 as lacking utility for the reasons set forth in the previous Office Action. The PTO relies on Pennica *et al.*, Haynes *et al.*, Hu *et al.*, and Chen *et al.* for the propositions that there is lack of correlation between DNA amplification and increased mRNA levels, and that polypeptide levels cannot be predicted from mRNA levels. The PTO asserts that “[g]iven the increase in amplified DNA (DNA copy number) for PRO1277” in normal esophagus and skin compared to esophageal and melanoma tumor “and the evidence provided by the current literature, it is clear that one skilled in the art would not assume that a higher expression of mRNA would correlate with increased polypeptide levels.” (Office Action at 5, emphasis added). The PTO argues that further research is required to determine the role of PRO1277 in cancer, making the asserted utility not substantial. The PTO also states that declarations submitted with Applicants’ previous response are insufficient to overcome the rejection.

#### **Utility – Legal Standard**

According to the Utility Examination Guidelines (“Utility Guidelines”), 66 Fed. Reg. 1092 (2001) an invention complies with the utility requirement of 35 U.S.C. § 101, if it has at least one asserted “specific, substantial, and credible utility” or a “well-established utility.”

Appl. No. : 10/063,540  
Filed : May 2, 2002

Under the Utility Guidelines, a utility is “specific” when it is particular to the subject matter claimed. For example, it is generally not enough to state that a nucleic acid is useful as a diagnostic tool without also identifying the condition that is to be diagnosed.

The requirement of “substantial utility” defines a “real world” use, and derives from the Supreme Court’s holding in *Brenner v. Manson*, 383 U.S. 519, 534 (1966) stating that “The basic *quid pro quo* contemplated by the Constitution and the Congress for granting a patent monopoly is the benefit derived by the public from an invention with substantial utility.” In explaining the “substantial utility” standard, M.P.E.P. § 2107.01 cautions, however, that Office personnel must be careful not to interpret the phrase “immediate benefit to the public” or similar formulations used in certain court decisions to mean that products or services based on the claimed invention must be “currently available” to the public in order to satisfy the utility requirement. “Rather, any reasonable use that an applicant has identified for the invention that can be viewed as providing a public benefit should be accepted as sufficient, at least with regard to defining a ‘substantial’ utility.” (M.P.E.P. § 2107.01, emphasis added).

The mere consideration that further experimentation might be performed to more fully develop the claimed subject matter does not support a finding of lack of utility. M.P.E.P. § 2107.01 III cites *In re Brana*, 51 F.3d 1560, 1566, 34 U.S.P.Q.2d 1436 (Fed. Cir. 1995) in stating that “Usefulness in patent law ... necessarily includes the expectation of further research and development. The stage at which an invention in this field becomes useful is well before it is ready to be administered to humans.” Further, “[T]o violate § 101 the claimed device must be totally incapable of achieving a useful result” *Juicy Whip Inc. v. Orange Bang Inc.*, 51 U.S.P.Q.2d 1700 (Fed. Cir. 1999), citing *Brooktree Corp. v. Advanced Micro Devices, Inc.*, 977 F.2d 1555, 1571 (Fed. Cir. 1992).

Indeed, the Guidelines for Examination of Applications for Compliance With the Utility Requirement, set forth in M.P.E.P. § 2107 II(B)(1) gives the following instruction to patent examiners: “If the applicant has asserted that the claimed invention is useful for any particular practical purpose ... and the assertion would be considered credible by a person of ordinary skill in the art, do not impose a rejection based on lack of utility.”

Finally, in assessing the credibility of the asserted utility, the M.P.E.P. states that “to overcome the presumption of truth that an assertion of utility by the applicant enjoys” the PTO

Appl. No. : 10/063,540  
Filed : May 2, 2002

must establish that it is “more likely than not that one of ordinary skill in the art would doubt (i.e., “question”) the truth of the statement of utility.” M.P.E.P. § 2107.02 III A. The M.P.E.P. cautions that:

Rejections under 35 U.S.C. 101 have been **rarely sustained** by federal courts. Generally speaking, **in these rare cases**, the 35 U.S.C. 101 rejection was sustained [] because the **applicant ... asserted a utility that could only be true if it violated a scientific principle, such as the second law of thermodynamics, or a law of nature, or was wholly inconsistent with contemporary knowledge in the art**. M.P.E.P. § 2107.02 III B., citing *In re Gazave*, 379 F.2d 973, 978, 154 U.S.P.Q. 92, 96 (CCPA 1967) (underline emphasis in original, bold emphasis added).

*Utility need NOT be Proved to a Statistical Certainty – a Reasonable Correlation between the Evidence and the Asserted Utility is Sufficient*

An Applicant's assertion of utility creates a presumption of utility that will be sufficient to satisfy the utility requirement of 35 U.S.C. § 101, “unless there is a reason for one skilled in the art to question the objective truth of the statement of utility or its scope.” *In re Langer*, 503 F.2d 1380, 1391, 183 USPQ 288, 297 (CCPA 1974). *See, also In re Jolles*, 628 F.2d 1322, 206 USPQ 885 (CCPA 1980); *In re Irons*, 340 F.2d 974, 144 USPQ 351 (1965); *In re Sichert*, 566 F.2d 1154, 1159, 196 USPQ 209, 212-13 (CCPA 1977). Compliance with 35 U.S.C. § 101 is a question of fact. *Raytheon v. Roper*, 724 F.2d 951, 956, 220 USPQ 592, 596 (Fed. Cir. 1983) cert. denied, 469 US 835 (1984). The evidentiary standard to be used throughout *ex parte* examination in setting forth a rejection is a preponderance of the evidence, or “more likely than not” standard. *In re Oetiker*, 977 F.2d 1443, 1445, 24 USPQ2d 1443, 1444 (Fed. Cir. 1992). This is stated explicitly in the M.P.E.P.:

[T]he applicant does not have to provide evidence sufficient to establish that an asserted utility is true “beyond a reasonable doubt.” **Nor must the applicant provide evidence such that it establishes an asserted utility as a matter of statistical certainty.** Instead, evidence will be sufficient if, considered as a whole, it leads a person of ordinary skill in the art to conclude that the asserted utility is more likely than not. M.P.E.P. at § 2107.02, part VII (2004) (underline emphasis in original, bold emphasis added, internal citations omitted).

The PTO has the initial burden to offer evidence “that one of ordinary skill in the art would reasonably doubt the asserted utility.” *In re Brana*, 51 F.3d 1560, 1566, 34 U.S.P.Q.2d 1436 (Fed. Cir. 1995). Only then does the burden shift to the Applicant to provide rebuttal

Appl. No. : 10/063,540  
Filed : May 2, 2002

evidence. *Id.* As stated in the M.P.E.P., such rebuttal evidence does not need to absolutely prove that the asserted utility is real. Rather, the evidence only needs to be reasonably indicative of the asserted utility.

In *Fujikawa v. Wattanasin*, 93 F.3d 1559, 39 U.S.P.Q.2d 1895 (Fed. Cir. 1996), the Court of Appeals for the Federal Circuit upheld a PTO decision that *in vitro* testing of a novel pharmaceutical compound was sufficient to establish practical utility, stating the following rule:

[T]esting is often required to establish practical utility. But the test results **need not absolutely prove** that the compound is pharmacologically active. All that is required is that the tests be “*reasonably* indicative of the desired [pharmacological] response.” In other words, there must be a **sufficient correlation** between the tests and an asserted pharmacological activity so as to convince those skilled in the art, **to a reasonable probability**, that the novel compound will exhibit the asserted pharmacological behavior.” *Fujikawa v. Wattanasin*, 93 F.3d 1559, 1564, 39 U.S.P.Q.2d 1895 (Fed. Cir. 1996) (internal citations omitted, bold emphasis added, italics in original).

While the *Fujikawa* case was in the context of utility for pharmaceutical compounds, the principals stated by the Court are applicable in the instant case where the asserted utility is for a therapeutic and diagnostic use – utility does not have to be established to an absolute certainty, rather, the evidence must convince a person of skill in the art “to a reasonable probability.” In addition, the evidence need not be direct, so long as there is a “sufficient correlation” between the tests performed and the asserted utility.

The Court in *Fujikawa* relied in part on its decision in *Cross v. Iizuka*, 753 F.2d 1040, 224 U.S.P.Q. 739 (Fed. Cir. 1985). In *Cross*, the Appellant argued that basic *in vitro* tests conducted in cellular fractions did not establish a practical utility for the claimed compounds. Appellant argued that more sophisticated *in vitro* tests using intact cells, or *in vivo* tests, were necessary to establish a practical utility. The Court in *Cross* rejected this argument, instead favoring the argument of the Appellee:

[I]n *vitro* results...are generally predictive of *in vivo* test results, i.e., there is a **reasonable correlation** therebetween. Were this not so, the testing procedures of the pharmaceutical industry would not be as they are. [Appellee] has not urged, and rightly so, that there is an invariable exact correlation between *in vitro* test results and *in vivo* test results. Rather, [Appellee's] position is that successful *in vitro* testing for a particular pharmacological activity establishes a **significant probability** that *in vivo* testing for this particular pharmacological activity will be

Appl. No. : 10/063,540  
Filed : May 2, 2002

successful. *Cross v. Iizuka*, 753 F.2d 1040, 1050, 224 U.S.P.Q. 739 (Fed. Cir. 1985) (emphasis added).

The *Cross* case is very similar to the present case. Like *in vitro* testing in the pharmaceutical industry, those of skill in the field of biotechnology rely on the reasonable correlation that exists between gene expression and protein expression (see below). Were there no reasonable correlation between the two, the techniques that measure gene levels such as microarray analysis, differential display, and quantitative PCR would not be so widely used by those in the art. As in *Cross*, Applicants here do not argue that there is “an invariable exact correlation” between gene expression and protein expression. Instead, Applicants’ position detailed below is that a measured change in gene expression in cancer cells establishes a “significant probability” that the expression of the encoded polypeptide in cancer will also be changed based on “a reasonable correlation therebetween.”

Taken together, the legal standard for demonstrating utility is a relatively low hurdle. An Applicant need only provide evidence such that it is **more likely than not that a person of skill in the art would be convinced, to a reasonable probability, that the asserted utility is true.** The evidence need not be direct evidence, so long as there is a reasonable correlation between the evidence and the asserted utility. The Applicant **does not need to provide evidence such that it establishes an asserted utility as a matter of statistical certainty.**

Even assuming that the PTO has met its initial burden to offer evidence that one of ordinary skill in the art would reasonably doubt the truth of the asserted utility, Applicants assert that they have met their burden of providing rebuttal evidence such that it is more likely than not those skilled in the art, to a reasonable probability, would believe that the claimed invention is useful as a diagnostic tool for cancer.

### **Substantial Utility**

#### **The Data in Example 18 are Data Regarding Differential mRNA Levels, not Gene Amplification**

Applicants begin by clarifying that the data concerning the differential expression of the PRO1277 gene presented in Example 18 relate to gene expression, not gene amplification. The description of Example 18 makes clear that the results were obtained by quantitative PCR amplification of cDNA libraries. It is well known in the art that cDNA libraries are made from

**Appl. No.** : 10/063,540  
**Filed** : May 2, 2002

mRNA, and reflect the level of mRNA for a particular gene in the source tissue. Thus, Example 18 is reporting a measure of the *expression* of the PRO1277 gene, i.e. mRNA levels, not its *amplification*, i.e. the number of copies of PRO1277 in the genome.

Throughout the Office Action, the PTO refers to Example 18 as “amplification data.” For example, the PTO states “[g]iven the increase in amplified DNA (DNA copy number) for PRO 1277” Office Action at 5 (emphasis added). Similarly, the PTO states that “the instant specification provides no information regarding differential mRNA levels of PRO1277 .... The specification describes only gene amplification data.” Office Action at 12 (emphasis added). In rejecting the declaration of Dr. Polakis, the PTO again states that “the instant specification provides no information regarding differential mRNA levels of PRO1277 .... Only gene amplification data were presented.” Office Action at 13-14 (emphasis added).

The PTO also continues to cite Pennica *et al.* to teach that “it does not necessarily follow that an increase in gene copy number results in increased gene expression.” Office Action at 16 (emphasis added). And although the PTO agrees that Pennica does not discuss the relationship between the mRNA level and the level of protein expression, the PTO states that Pennica was cited “to show the lack of correlation of between DNA amplification and gene expression.” Office Action at 16 (emphasis in original). Elsewhere the PTO states that “one cannot determine from the data in the specification whether the observed ‘amplification’ of nucleic acid is due to increase in copy number, or alternatively due to increase in transcription rates.” Office Action at 12.

Whether or not gene amplification leads to increased gene expression is irrelevant to this particular application. Likewise, whether the differential mRNA expression of the PRO1277 gene reported in Example 18 is due to an increase or decrease in copy number, or alternatively due to an increase or decrease in transcription rates is simply not relevant. Applicants have provided reliable evidence that the PRO1277 mRNA is differentially expressed in certain tumors. Whether this differential expression is due to changes in gene copy number, transcription rates, a combination of the two, or some other known or unknown cellular mechanism is simply not relevant to Applicants’ asserted utility.

Appl. No. : 10/063,540  
Filed : May 2, 2002

Summary of Applicants' Arguments and the PTO's Response

In an attempt to clarify Applicants' argument, Applicants offer a summary of their argument and the disputed issues involved. Applicants assert that the claimed antibodies have utility as diagnostic tools for cancer, particularly esophageal and melanoma cancer. Applicants' asserted utility rests on the following argument:

1. Applicants have provided reliable evidence that mRNA for the PRO1277 polypeptide is expressed at least two-fold higher in normal esophageal and skin tissue than compared to esophageal and melanoma tumor, respectively;

2. Applicants assert that it is well-established in the art that a change in the level of mRNA for a particular protein, e.g. an increase, generally leads to a corresponding change in the level of the encoded protein, e.g. an increase;

3. Given Applicants' evidence that the level of mRNA for the PRO1277 polypeptide is decreased in esophageal and melanoma tumors compared to normal esophageal and skin tissue, respectively, it is likely that the PRO1277 polypeptide is expressed at decreased levels in esophageal and melanoma tumors compared to normal esophageal and skin tissue, respectively;

4. Antibodies to proteins which are differentially expressed in certain tumors are useful as diagnostic tools.

Applicants understand the PTO to be making several arguments in response to Applicants' asserted utility:

1. The PTO has challenged the reliability of the evidence reported in Example 18, and states that it provides no information regarding differential mRNA expression;

2. The PTO cites Pennica *et al.* to support its position that gene amplification is not necessarily correlated to gene expression;

3. The PTO cites Hu *et al.* Haynes *et al.* and Chen *et al.* to support its assertion that the literature cautions against drawing conclusions based on small changes in transcript expression levels between normal and cancerous tissue, and that mRNA levels are not predictive of protein levels;

4. The PTO concludes that based on the cited literature, one of skill in the art would not assume that higher expression of mRNA would correlate with increased polypeptide levels.



Appl. No. : 10/063,540  
Filed : May 2, 2002

Therefore, further research needs to be done to determine if the increase or decrease in PRO1277 DNA supports a role for the peptide in cancerous tissue. (See Office Action at 5-6).

As detailed below, Applicants submit that the PTO has failed to demonstrate that this is one of the “rare cases” where the applicants have “asserted a utility that could only be true if it violated a scientific principle, such as the second law of thermodynamics, or a law of nature, or was wholly inconsistent with contemporary knowledge in the art.” M.P.E.P. § 2107.02 III B. First, the PTO has failed to offer any evidence to support its rejection of the data in Example 18 and the Declaration of Chris Grimaldi in support of these data. Second, as discussed above and can be seen from Applicants’ summary of their argument, Applicants submit that any lack of correlation between gene amplification and gene expression is not at issue in this application and therefore the Pennica *et al.* reference is not relevant. Third, Applicants submit that the Haynes *et al.* Hu *et al.* and Chen *et al.* references are not contrary to Applicants’ arguments, and therefore are not evidence to support the PTO’s position. Finally, even if the PTO has met its initial burden, Applicants have submitted enough rebuttal evidence such that it is **more likely than not** that a person of skill in the art would be convinced, **to a reasonable probability**, that the asserted utility is true. As stated above, Applicants’ evidence need not be direct evidence, so long as there is a reasonable correlation between the evidence and the asserted utility. **The standard is not absolute certainty.**

*Applicants have established that the Gene Encoding the PRO1277 Polypeptide is Differentially Expressed in Certain Cancers compared to Normal Tissue*

Applicants first address the PTO’s previous argument the evidence of differential expression of the gene encoding the PRO1277 polypeptide in esophageal and melanoma tumors is insufficient because it does not teach what the normal level of expression is, it does not indicate how high the expression level is compared to the disease tissue, it lacks statistical correlation, there is no data to compare expression in the normal and disease samples, and that because the normal and tumor samples are not from the same person, there is no possibility of direct comparison between the normal and tumor samples. Applicants submit that the gene expression data provided in Example 18 of the present application are sufficient to establish that the PRO1277 gene is differentially expressed in esophageal and melanoma tumors.

Appl. No. : 10/063,540  
Filed : May 2, 2002

The gene expression data in the specification, Example 18, shows that the mRNA associated with protein PRO1277 was more highly expressed in normal esophageal and skin tissue versus esophageal and melanoma tumor. Gene expression was analyzed using standard semi-quantitative PCR amplification reactions of cDNA libraries isolated from different human tumor and normal human tissue samples. Identification of the differential expression of the PRO1277 polypeptide-encoding gene in tumor tissue compared to the corresponding normal tissue renders the molecule useful as a diagnostic tool for the determination of the presence or absence of tumor. Applicants previously submitted a first Declaration of J. Christopher Grimaldi, an expert in the field of cancer biology. This declaration explains the importance of the data in Example 18, and how differential gene and protein expression studies are used to differentiate between normal and tumor tissue (see Declaration, paragraph 7).

In paragraph 5 of his declaration, Mr. Grimaldi states that the gene expression studies reported in Example 18 of the instant application were made from pooled samples of normal and of tumor tissues. Contrary to the PTO's assertions that this makes the data unreliable, Mr. Grimaldi explains that:

The DNA libraries used in the gene expression studies were made from pooled samples of normal and of tumor tissues. *Data from pooled samples is more likely to be accurate than data obtained from a sample from a single individual.* That is, the detection of variations in gene expression is likely to represent a more generally relevant condition when pooled samples from normal tissues are compared with pooled samples from tumors in the same tissue type. (Paragraph 5) (emphasis added).

In paragraphs 6 and 7, Mr. Grimaldi explains that the semi-quantitative analysis employed to generate the data of Example 18 is sufficient to determine if a gene is over- or under-expressed in tumor cells compared to corresponding normal tissue. He states that any visually detectable difference seen between two samples is indicative of at least a two-fold difference in cDNA between the tumor tissue and the counterpart normal tissue. Thus, the results of Example 18 reflect at least a two-fold difference between normal and tumor samples. He also states that the results of the gene expression studies indicate that the genes of interest "can be used to differentiate tumor from normal," thus establishing their reliability. He explains that, contrary to the PTO's assertions, "The precise levels of gene expression are irrelevant; what matters is that there is a relative difference in expression between normal tissue and tumor

**Appl. No.** : **10/063,540**  
**Filed** : **May 2, 2002**

tissue.” (Paragraph 7). Thus, since it is the relative level of expression between normal tissue and suspected cancerous tissue that is important, the precise level of expression in normal tissue is irrelevant. Likewise, there is no need for quantitative data to compare the level of expression in normal and tumor tissue. As Mr. Grimaldi states, “If a difference is detected, this indicates that the gene and its corresponding polypeptide and antibodies against the polypeptide are useful for diagnostic purposes, to screen samples to differentiate between normal and tumor.”

The PTO has stated that the Grimaldi Declaration is insufficient to overcome the rejection of Claims 1-5, offering two arguments. First, the PTO argues that “there is no evidentiary art that would corroborate for example, that ‘any visually detectable difference seen between two samples is indicative of at least a two-fold difference in cDNA between the tumor tissue and the counterpart normal tissue.’” Office Action at 13 (quoting first Grimaldi Declaration).

Applicants submit that the declaration of Mr. Grimaldi is based on personal knowledge of the relevant facts at issue. Mr. Grimaldi is an expert in the field and conducted or supervised the experiments at issue. Applicants remind the PTO that “[o]ffice personnel must accept an opinion from a qualified expert that is based upon relevant facts whose accuracy is not being questioned.” PTO Utility Examination Guidelines (2001) (emphasis added). In addition, declarations relating to issues of fact should not be summarily dismissed as “opinions” without an adequate explanation of how the declaration fails to rebut the Examiner’s position. *In re Alton* 76 F.3d 1168 (Fed. Cir. 1996).

In addition, Applicants provide herewith as Exhibit 1 a copy of page 122 of the 2002-2003 New England Biolabs catalog. Exhibit 1 shows DNA size markers of differing lengths run on an agarose gel. The column on the left provides the mass of each marker in nanograms and the column on the right provides the length of the marker. It is apparent that the band intensity of markers having mass differences of two-fold are readily distinguishable by eye (*see e.g.*, the difference in band intensities of the 0.1kb fragment present at 61ng and the 0.5kb marker present at 124ng). Accordingly, Applicants maintain that the procedures used to detect differences in expression levels were sufficiently sensitive to detect two-fold differences.

The PTO has not supplied any reasons or evidence to question the accuracy of the facts upon which Mr. Grimaldi based his opinion. Mr. Grimaldi has personal knowledge of the relevant facts, has based his opinion on those facts, and the PTO has offered no reason or

**Appl. No.** : 10/063,540  
**Filed** : May 2, 2002

evidence to reject either the underlying facts or his opinion. Therefore, the PTO should accept Mr. Grimaldi's opinion with regard to his statement that "any visually detectable difference seen between two samples is indicative of at least a two-fold difference in cDNA between the tumor tissue and the counterpart normal tissue" and that the nucleic acids of interest "can be used to differentiate tumor from normal." Together, these statements establish that there is at least a two-fold difference in expression, and that the results are reliable enough that they can be used to distinguish tumor from normal tissue.

Second, the PTO states that with regard to the Grimaldi declaration, "one cannot determine from the data in the specification whether the observed 'amplification' of nucleic acid is due to increase in copy number, or alternatively due to increase in transcription rates," and that the specification does not provide any information regarding "differential mRNA levels of PRO1277," but rather "describes only gene amplification." Office Action at 12.

Applicant's again point out that the data in Example 18 are gene expression data, not gene amplification data. The specification and the first Grimaldi Declaration make clear that Example 18 used semi-quantitative PCR of cDNA libraries. Therefore, one of skill in the art would know that Example 18 is a measure of mRNA levels, and reflects differential PRO1277 gene expression, not gene amplification. As discussed above, the PTO cites Pennica *et al.* for support of the argument that gene amplification does not necessarily lead to increased gene expression. This reference is irrelevant to the instant application which reports differential gene expression, not gene amplification. Therefore, this reference and the PTO's arguments regarding gene amplification do not support the PTO's challenge of the sufficiency of the Example 18 data, or the first Grimaldi Declaration.

The PTO also cites Hu *et al.* (J. Proteome Res., 2(4):405-12 (2003)) for support for its assertion the literature cautions researchers from drawing conclusions based on small changes in transcript expression levels between normal and cancerous tissue. The PTO states that Hu teaches that not all genes with increased expression in cancer have a known or published role in cancer.

In Hu, the researchers used an automated literature-mining tool to summarize and estimate the relative strengths of all human gene-disease relationships published on Medline. They then generated a microarray expression dataset comparing breast cancer and normal breast

Appl. No. : 10/063,540  
Filed : May 2, 2002

tissue. Using their data-mining tool, they looked for a correlation between the strength of the literature association between the gene and breast cancer, and the magnitude of the difference in expression level. They report that for genes displaying a 5-fold change or less in tumors compared to normal, there was no evidence of a correlation between altered gene expression and a *known* role in the disease. *See* Hu at 411. However, among genes with a 10-fold or more change in expression level, there was a strong correlation between expression level and a *published* role in the disease. *Id.* at 412. Importantly, Hu reports that the observed correlation was only found among estrogen receptor-positive tumors, not ER-negative tumors. *Id.*

The general findings of Hu are not surprising – one would expect that genes with the greatest change in expression in a disease would be the first targets of research, and therefore have the strongest known relationship to the disease as measured by the number of publications reporting a connection with the disease. The correlation reported in Hu only indicates that the greater the change in expression level, the more likely it is that there is a *published* or *known* role for the gene in the disease, as found by their automated literature-mining software. Thus, Hu's results merely reflect a bias in the literature toward studying the most prominent targets, and reflect nothing regarding the ability of a gene that is 2-fold or more differentially expressed in tumors to serve as a disease marker.

Hu acknowledges the shortcomings of this method in explaining the disparity in Hu's findings for ER-negative versus ER-positive tumors: Hu attributes the "bias in the literature" toward the more prevalent ER-positive tumors as the explanation for the lack of any correlation between number of publications and gene expression levels in less-prevalent (and, therefore, less studied) ER-negative tumors. *Id.* Because of this intrinsic bias, Hu's methodology is unlikely to ever note a correlation of a disease with less differentially-expressed genes and their corresponding proteins, regardless of whether or not an actual relationship between the disease and less differentially-expressed genes exists. Accordingly, Hu's methodology yields results that provide little or no information regarding biological significance of genes with less than 5-fold expression change in disease. Nowhere in Hu does it say that a lack of correlation in their study means that genes with a less than five-fold change in level of expression in cancer cannot serve as a molecular marker of cancer.

**Appl. No.** : 10/063,540  
**Filed** : May 2, 2002

Applicants submit that a lack of known role for PRO1277 in cancer does not prevent its use as a diagnostic tool for cancer. There is a difference between use of a gene for distinguishing between tumor and normal tissue on the one hand, and establishing a role for the gene in cancer on the other. Genes with lower levels of change in expression may or may not be the most important genes in causing the disease, but the genes can still show a consistent and measurable change in expression. While such genes may or may not be good targets for further research, they can nonetheless be used as diagnostic tools. Thus, Hu does not refute the Applicants' assertion that the PRO1277 gene can be used as a cancer diagnostic tool because it is differentially expressed in certain tumors.

Contrary to the PTO's assertion that one must know what role a gene or polypeptide plays in cancer for it have utility, the PTO's own written policies recognize that the utility of a nucleic acid does not depend on the function of the encoded gene product. The Utility Examination Guidelines published on January 5, 2001 state: "In addition, the utility of a claimed DNA does not necessarily depend on the function of the encoded gene product. A claimed DNA may have a specific and substantial utility because, e.g. it hybridizes near a disease-associated gene or it has a gene regulating activity." (Federal Register, Volume 66, page 1095, Comment 14); *see also Exhibits 2-4* (U.S. Patent Nos. 6,465,185, 6,228,582, and 6,162,604) (patents on polymorphisms which are indicative of a predisposition to a particular condition are patentable even though they may or may not cause the disease itself). Similarly, here the disclosed nucleic acids, as well as the encoded polypeptides and related antibodies, are useful for determining whether an individual has cancer regardless of whether or not they are the cause of the cancer.

The position of the PTO requiring a known role for PRO1277 in cancer for utility is also inconsistent with the analogous standard for therapeutic utility of a compound where "the mere identification of a pharmacological activity of a compound that is relevant to an asserted pharmacological use provides an 'immediate benefit to the public' and thus satisfies the utility requirement." M.P.E.P. §2701.01 (emphasis original). Here, the mere identification of altered expression in tumors is relevant to diagnosis of tumors, and, therefore, provides an immediate benefit to the public.

In conclusion, Applicants submit that the evidence reported in Example 18, combined with the first Grimaldi Declaration, establish that there is at least a two-fold difference in

Appl. No. : 10/063,540  
Filed : May 2, 2002

PRO1277 cDNA between normal esophageal and skin tissue compared to esophageal and melanoma tumor, respectively. Therefore, it follows that expression levels of the PRO1277 gene can be used to distinguish esophageal and melanoma tumor tissue from normal esophageal and skin tissue, respectively. The PTO has not offered any significant arguments or evidence to the contrary. As Applicants explain below, it is more likely than not that the PRO1277 polypeptide can also be used to distinguish esophageal and melanoma tumor tissue from normal esophageal and skin tissue, respectively. This provides utility for the claimed antibodies to the PRO1277 polypeptides.

*Applicants have established that the Accepted Understanding in the Art is that there is a Positive Correlation between mRNA Levels and the Level of Expression of the Encoded Protein*

Applicants next turn to the second portion of their argument in support of their asserted utility – that it is well-established in the art that a change in the level of mRNA for a particular protein generally leads to a corresponding change in the level of the encoded protein; given Applicants’ evidence of differential expression of the mRNA for the PRO1277 polypeptide in esophageal and melanoma tumor, it is likely that the PRO1277 polypeptide is differentially expressed; and antibodies to proteins differentially expressed in certain tumors have utility as diagnostic tools.

In response to Applicants’ assertion, the PTO cites Haynes *et al.* (Electrophoresis, 19(11):1862-71 (1998)) and Chen *et al.* (Mol. and Cell. Proteomics, 1:304-313 (2002)) as support for its argument that “polypeptide levels cannot be accurately predicted from mRNA levels.” Office Action at 16. For the reasons discussed below, Haynes is not contrary to Applicants’ asserted utility.

Haynes studied whether there is a correlation between the level of mRNA expression and the level of protein expression for 80 selected genes from yeast. The genes were selected because they constituted a relatively homogeneous group with respect to predicted half-life and expression level of the protein products. *See* Haynes at 1863. Haynes did not examine whether a change in transcript level for a particular gene led to a change in the level of expression of the corresponding protein. Instead, Haynes determined whether the steady-state transcript level

Appl. No. : 10/063,540  
Filed : May 2, 2002

correlated with the steady-state level of the corresponding protein based on an analysis of 80 different genes.

Haynes reported to have “found a general trend but no strong correlation between protein and transcript levels.” *Id.* However, a cursory inspection of Fig. 1 shows a clear correlation between the mRNA levels and protein levels measured. This correlation is confirmed by an inspection of the full-length research paper from which the data in Fig. 1 were derived, presented herein as Exhibit 5 (Gygi *et al.*, Molecular and Cellular Biology, Mar. 1999, 1720-1730). Gygi states that “there was a general trend of increased protein levels resulting from increased mRNA levels,” with a correlation coefficient of 0.935, indicating a strong correlation. Gygi at 1726. Moreover, Gygi also states that the correlation is especially strong for highly expressed mRNAs. *Id.* Thus, it is not clear that Haynes even supports the Examiner’s position, as Haynes did report a general trend, and Gygi reports a strong correlation between increasing mRNA levels and increasing protein levels.

The PTO focuses on the portion of Haynes where the authors reported that for some of the studied genes with equivalent mRNA levels, there were differences in corresponding protein expression, including some that varied by more than 50-fold. Similarly, Haynes reports that different proteins with similar expression levels were maintained by transcript levels that varied by as much as 40-fold. *Id.* Thus, Haynes showed that for one type of yeast, similar mRNA levels for *different* genes did not universally result in equivalent protein levels for the *different* gene products, and similar protein levels for *different* gene products did not universally result from equivalent mRNA levels for the *different* genes. These results are expected, since there are many factors that determine translation efficiency for a given transcript, or the half-life of the encoded protein. Not surprisingly, based on these results, Haynes concluded that protein levels cannot always be accurately predicted from the level of the corresponding mRNA transcript *when looking at the level of transcripts across different genes.*

Importantly, Haynes did not say that for a single gene, the level of mRNA transcript is not positively correlated with the level of protein expression. Applicants have asserted that increasing or decreasing the level of mRNA for the same gene leads to a increase or decrease for the corresponding protein. Haynes did not study this issue and says absolutely nothing about it.



**Appl. No.** : **10/063,540**  
**Filed** : **May 2, 2002**

Therefore, Haynes is not inconsistent with or contradictory to the utility of the instant claims, and offers no support for the PTO's position.

The PTO also cites Chen *et al.* for support for the assertion that polypeptide levels cannot be accurately predicted from mRNA levels. In Chen, the authors examined the relationship between mRNA levels and protein levels in 76 lung adenocarcinomas and 9 non-tumor lung samples. Like Haynes, Chen examined the global relationship between mRNA and the corresponding protein abundance by calculating the average mRNA and protein level of all the samples for each gene or protein, and then looked for a correlation across different genes. As discussed above with respect to Haynes, this measurement of a correlation across genes is not relevant to Applicants' asserted utility. Chen also looked at the level of mRNA of 98 individual genes and their corresponding proteins across the samples. Chen reports that 21.4% (21 of 98) of the genes showed a statistically significant correlation between protein and mRNA expression.

Chen provides scant evidence to counter Applicants' asserted utility for the claimed antibodies because portions of Chen support Applicants' assertions, and the remaining portions provide little insight into the relationship between mRNA levels and corresponding protein levels for mRNA that is differentially expressed in tumor cells relative to normal cells. Rather than looking for mRNAs which were differentially expressed, Chen merely selected proteins whose identity could be determined regardless of any changes in expression level (Chen at 306, right column). Importantly, it is not known if there was any substantial difference in mRNA levels for the various genes across samples – in short, with the exception of the genes in Figures 2A-2C, it is not known if the genes examined were differentially expressed. Also of significance for Applicants' asserted utility is the fact that Chen did not attempt to examine any differential expression between the cancerous lung samples and the non-cancerous lung samples – Chen did not distinguish between cancer and normal samples in their analysis.

Applicants have asserted that changes in mRNA levels, particularly those which are two-fold or greater, will correspond with measurable changes in polypeptide expression. The data in Chen support Applicants' assertion. In Figures 2A-2C, Chen plots mRNA value vs. protein value for three genes. In these figures, a wide range of mRNA expression levels were observed (approximately seven- to eight-fold), and a correlation between mRNA and protein levels was observed for all three mRNA/protein pairs. This supports Applicants' asserted correlation

**Appl. No.** : **10/063,540**  
**Filed** : **May 2, 2002**

between changes in mRNA levels which are two-fold or greater and changes in polypeptide expression.

The PTO relies on the fact that Chen also reports a lack of correlation for some mRNA/protein pairs to support its assertion that polypeptide levels cannot be accurately predicted from mRNA levels. However, the lack of correlation reported by Chen could be a result of a lack of substantial changes in mRNA level. This can be understood by again turning to Figures 2A-2C. As noted above, where a wide range of mRNA expression levels are seen, a correlation between mRNA and protein levels was observed. However, if one examines the data points within a small range of mRNA levels for these same genes, e.g. 500-600 or 5000-6000 in Figs. 2A-2C, it is clear that a correlation would not be detected for the data within this range. This does not mean that a correlation between changes in mRNA and protein does not exist for these genes, as is evident when larger changes in mRNA expression are included in the analysis. Instead, this indicates that for relatively small changes in mRNA, any correlation is masked by imprecision in the measurements.

Chen's experiment compared mRNA levels vs. protein levels across samples without selecting mRNA that showed a difference in expression level. And unlike Applicants, Chen did not examine differences in mRNA between tumor and normal tissue. Since almost all samples tested by Chen were from the same type of tissue, few substantial variations in the level of mRNA or protein for a particular gene across the samples tested would be expected. Instead, it would be expected that most genes examined by Chen would have similar mRNA or protein levels across the samples. Figures 2A-2C of Chen demonstrate that the methods utilized by Chen cannot detect correlations between mRNA and protein levels when only small differences in mRNA expression are observed, but a correlation is detected when larger differences in mRNA expression are observed.

Accordingly, the only data reported by Chen which shows substantial changes in the expression of mRNA, Figures 2A-C, confirms Applicants' assertion that substantial changes in mRNA levels (e.g., 2-fold or greater) will correspond to substantial changes in polypeptide expression. Further, this data also explains the lack of observed correlation between mRNA levels and protein levels for other genes reported by Chen. Thus, even given Chen's inability to

**Appl. No.** : 10/063,540  
**Filed** : May 2, 2002

detect a correlation between mRNA and protein in some genes, Chen's results do not refute Applicants' position.

Instead, Chen supports Applicants' position that a significant correlation between mRNA and protein levels exists for changes in mRNA levels that are 2-fold or greater. In further support of Applicants' position, Chen cites Celis *et al.* (FEBS Lett., 480:2-16 (2000)) stating that the authors "found a good correlation between transcript and protein levels among 40 well resolved, abundant proteins using a proteomic and microarray study of bladder cancer." *Chen* at 311, first column (emphasis added). As mentioned above, the lack of a correlation across genes is not relevant to Applicants' asserted utility, and therefore Chen's discussion of this issue and citation of Anderson and Seilhamer (Electrophoresis, 18:533-37 (1997)) and Gygi *et al.* (Mol. Cell. Bio., 19:1720-30 (1999)) offer no support for the PTO's position.

Even if the results in Chen supported the PTO's argument, which they do not as discussed above, one contrary example does not establish that one of skill in the art would find it is more likely than not there is no general correlation between changes in mRNA level and changes in protein level for an individual gene. There are other non-transcriptional mechanisms for regulating gene and protein expression (*i.e.*, post-transcriptional regulation of genes, translation efficiency, etc.). However, as shown by the declarations, references, and textbooks discussed below, Applicants submit that the understanding in the art is that generally there is a correlation between a change in mRNA level and a change in protein level. In fact, the working hypothesis among those skilled in the art, as illustrated by the evidence presented by Applicants, is that there is a positive correlation between changes in mRNA levels and changes in protein levels for a particular gene.

In support of the assertion that changes in mRNA are positively correlated to changes in protein levels, Applicants previously submitted a copy of a second Declaration by J. Christopher Grimaldi, an expert in the field of cancer biology. As stated in paragraph 5 of the declaration, "Those who work in this field are well aware that in the vast majority of cases, when a gene is over-expressed...the gene product or polypeptide will also be over-expressed.... This same principal applies to gene under-expression." Further, "the detection of increased mRNA expression is expected to result in increased polypeptide expression, and the detection of decreased mRNA expression is expected to result in decreased polypeptide expression. The

Appl. No. : 10/063,540  
Filed : May 2, 2002

detection of increased or decreased polypeptide expression can be used for cancer diagnosis and treatment.” The references cited in the declaration and submitted herewith support this statement.

Applicants also previously submitted a copy of the declaration of Paul Polakis, Ph.D., an expert in the field of cancer biology. As stated in paragraph 6 of his declaration:

Based on my own experience accumulated in more than 20 years of research, including the data discussed in paragraphs 4 and 5 above [showing a positive correlation between mRNA levels and encoded protein levels in the vast majority of cases] and my knowledge of the relevant scientific literature, it is my considered scientific opinion that for human genes, an increased level of mRNA in a tumor cell relative to a normal cell typically correlates to a similar increase in abundance of the encoded protein in the tumor cell relative to the normal cell. In fact, it remains a central dogma in molecular biology that increased mRNA levels are predictive of corresponding increased levels of the encoded protein. (Emphasis added).

Dr. Polakis acknowledges that there are published cases where such a correlation does not exist, but states that it is his opinion, based on over 20 years of scientific research, that “such reports are exceptions to the commonly understood general rule that increased mRNA levels are predictive of corresponding increased levels of the encoded protein.” (Polakis Declaration, paragraph 6).

The statements of Grimaldi and Polakis are supported by the teachings in Molecular Biology of the Cell, a leading textbook in the field (Bruce Alberts, *et al.*, Molecular Biology of the Cell (3<sup>rd</sup> ed. 1994) (submitted herewith as Exhibit 6) and (4<sup>th</sup> ed. 2002) (submitted herewith as Exhibit 7)). Figure 9-2 of Exhibit 6 shows the steps at which eukaryotic gene expression can be controlled. The first step depicted is transcriptional control. Exhibit 6 provides that “[f]or most genes transcriptional controls are paramount. This makes sense because, of all the possible control points illustrated in Figure 9-2, only transcriptional control ensures that no superfluous intermediates are synthesized.” Exhibit 6 at 403 (emphasis added). In addition, the text states that “Although controls on the initiation of gene transcription are the predominant form of regulation for most genes, other controls can act later in the pathway from RNA to protein to modulate the amount of gene product that is made.” Exhibit 6 at 453 (emphasis added). Thus, as established in Exhibit 6, the predominant mechanism for regulating the amount of protein produced is by regulating transcription initiation.

Appl. No. : 10/063,540  
Filed : May 2, 2002

In Exhibit 7, Figure 6-3 on page 302 illustrates the basic principle that there is a correlation between increased gene expression and increased protein expression. The accompanying text states that “a cell can change (or regulate) the expression of each of its genes according to the needs of the moment – *most obviously by controlling the production of its mRNA.*” Exhibit 7 at 302 (emphasis added). Similarly, Figure 6-90 on page 364 of Exhibit 7 illustrates the path from gene to protein. The accompanying text states that while potentially each step can be regulated by the cell, “the initiation of transcription is the most common point for a cell to regulate the expression of each of its genes.” Exhibit 7 at 364 (emphasis added). This point is repeated on page 379, where the authors state that of all the possible points for regulating protein expression, “[f]or most genes transcriptional controls are paramount.” Exhibit 7 at 379 (emphasis added).

Further support for Applicants’ position can be found in the textbook, *Genes VI*, (Benjamin Lewin, *Genes VI* (1997)) (submitted herewith as Exhibit 8) which states “having acknowledged that control of gene expression can occur at multiple stages, and that production of RNA cannot inevitably be equated with production of protein, it is clear that the overwhelming majority of regulatory events occur at the initiation of transcription.” *Genes VI* at 847-848 (emphasis added).

Additional support is also found in Zhigang *et al.*, *World Journal of Surgical Oncology* 2:13, 2004, submitted herewith as Exhibit 9. Zhigang studied the expression of prostate stem cell antigen (PSCA) protein and mRNA to validate it as a potential molecular target for diagnosis and treatment of human prostate cancer. The data showed “a high degree of correlation between PSCA protein and mRNA expression” Exhibit 9 at 4. Of the samples tested, 81 out of 87 showed a high degree of correlation between mRNA expression and protein expression. The authors conclude that “it is demonstrated that PSCA protein and mRNA overexpressed in human prostate cancer, and that the increased protein level of PSCA was resulted from the upregulated transcription of its mRNA.” Exhibit 9 at 6. Even though the correlation between mRNA expression and protein expression occurred in 93% of the samples tested, not 100%, the authors state that “PSCA may be a promising molecular marker for the clinical prognosis of human Pca and a valuable target for diagnosis and therapy of this tumor.” Exhibit 9 at 7.

Appl. No. : 10/063,540  
Filed : May 2, 2002

Further, Meric *et al.*, Molecular Cancer Therapeutics, vol. 1, 971-979 (2002), submitted herewith as Exhibit 10, states the following:

The **fundamental principle** of molecular therapeutics in cancer is to exploit the differences in gene expression between cancer cells and normal cells...[M]ost efforts have concentrated on identifying differences in gene expression at the level of mRNA, which can be attributable to either DNA amplification or to differences in transcription. Meric *et al.* at 971 (emphasis added).

Those of skill in the art would not be focusing on differences in gene expression between cancer cells and normal cells if there were no correlation between gene expression and protein expression.

Together, the declarations of Grimaldi and Polakis, the accompanying references, and the excerpts and references provided above all establish that the accepted understanding in the art is that there is a reasonable correlation between changes in gene expression and the level of the encoded protein.

In response to the second Grimaldi Declaration and the Polakis Declaration, the PTO states that “it is important to note that the instant specification provides no information regarding differential mRNA levels of PRO1277.... Only gene amplification data were presented. Therefore, the declaration is insufficient to overcome the rejection of claims 1-5 ... since it is limited to a discussion of data regarding the correlation of mRNA levels and polypeptides.” Office Action at 13-14. Applicants again reiterate that Example 18 is information regarding differential mRNA levels of PRO1277. Thus, the PTO’s rejection of the second Grimaldi Declaration and Polakis Declaration because they discuss the correlation of mRNA levels and polypeptide levels is misplaced.

The PTO also argues that further research needs to be done to determine whether the increase or decrease of PRO1277 DNA supports a role for the peptide in the cancerous tissue, and that such a role has not been suggested by the instant disclosure. *See* Office Action at 5-6.

Applicants submit that a lack of known role for PRO1277 in cancer does not prevent its use as a diagnostic tool for cancer. The fact that there is no known translocation or mutation of PRO1277, for example, is irrelevant to whether its differential expression can be used to assist in diagnosis of cancer – one does not need to know why PRO1277 is differentially expressed, or what the consequence of the differential expression is, in order to exploit the differential

Appl. No. : 10/063,540  
Filed : May 2, 2002

expression to distinguish tumor from normal tissue. In fact the Revised Interim Utility Guidelines promulgated by the PTO recognize that proteins which are differentially expressed in cancer have utility. (See the caveat in Example 12 which state that the utility requirement is satisfied where a protein is expressed in melanoma cells but not on normal skin and antibodies against the protein can be used to diagnose cancer.) In addition, while Applicants appreciate that actions taken in other applications are not binding on the PTO with respect to the present application, Applicants note that the PTO has issued several patents claiming differentially expressed polypeptides and antibodies to the same, or methods employing such antibodies. (See, e.g., U.S. Patent No. 6,414,117, U.S. Patent No. 6,124,433, U.S. Patent No. 6,156,500, and U.S. Patent No. 6,562,343 attached hereto as Exhibits 11-14.)

Accordingly, Applicants submit that they have offered sufficient evidence to establish that it is more likely than not that one of skill in the art would believe that because the PRO1277 mRNA is more highly expressed in normal esophageal and skin tissue than compared to esophageal and melanoma tumor, respectively, the PRO1277 polypeptide will also be more highly expressed in normal esophageal and skin tissue than compared to esophageal and melanoma tumor, respectively. This differential expression of the PRO1277 polypeptide makes antibodies to it useful as diagnostic tools for cancer.

*The Arguments made by the PTO are Not Sufficient to satisfy the PTO's Initial Burden of Offering Evidence "that one of ordinary skill in the art would reasonably doubt the asserted utility"*

As stated above, an Applicant's assertion of utility creates a presumption of utility that will be sufficient to satisfy the utility requirement of 35 U.S.C. § 101, "unless there is a reason for one skilled in the art to question the objective truth of the statement of utility or its scope." *In re Langer*, 503 F.2d 1380, 1391, 183 USPQ 288, 297 (CCPA 1974). The evidentiary standard to be used throughout *ex parte* examination in setting forth a rejection is a preponderance of the evidence, or "more likely than not" standard. *In re Oetiker*, 977 F.2d 1443, 1445, 24 USPQ2d 1443, 1444 (Fed. Cir. 1992). This is stated explicitly in the M.P.E.P.:

[T]he applicant does not have to provide evidence sufficient to establish that an asserted utility is true "beyond a reasonable doubt." **Nor must the applicant provide evidence such that it establishes an asserted utility as a matter of**

Appl. No. : 10/063,540  
Filed : May 2, 2002

**statistical certainty.** Instead, evidence will be sufficient if, considered as a whole, it leads a person of ordinary skill in the art to conclude that the asserted utility is more likely than not. M.P.E.P. at § 2107.02, part VII (2004) (underline emphasis in original, bold emphasis added, internal citations omitted).

The PTO has the initial burden to offer evidence “that one of ordinary skill in the art would reasonably doubt the asserted utility.” *In re Brana*, 51 F.3d 1560, 1566, 34 U.S.P.Q.2d 1436 (Fed. Cir. 1995). Only then does the burden shift to the Applicant to provide rebuttal evidence. *Id.* As stated in the M.P.E.P., such rebuttal evidence does not need to absolutely prove that the asserted utility is real. Rather, the evidence only needs to be reasonably indicative of the asserted utility.

Applicants remind the PTO that the M.P.E.P. cautions that rejections for lack of utility are rarely sustained by federal courts, and that generally speaking, a utility rejection was sustained because the applicant asserted a utility “that could **only be true if it violated a scientific principle, such as the second law of thermodynamics, or a law of nature, or was wholly inconsistent with contemporary knowledge in the art.**” M.P.E.P. § 2107.02 III B., citing *In re Gazave*, 379 F.2d 973, 978, 154 U.S.P.Q. 92, 96 (CCPA 1967) (underline emphasis in original, bold emphasis added). Rather than being wholly inconsistent with contemporary knowledge in the art, Applicants’ asserted utility is squarely within the teaching of leading textbooks in the field, and is supported by references and the declarations of skilled experts.

The PTO has not offered any arguments or cited any references to establish “that one of ordinary skill in the art would reasonably doubt” that antibodies to a polypeptide differentially expressed in certain tumors can be used as a diagnostic tool. Pennica *et al.* is irrelevant to the utility of the claimed antibodies as Example 18 reports gene expression data, not gene amplification data. Likewise, neither Hu *et al.* nor Haynes *et al.* supports the PTO’s position or is contrary to Applicants’ asserted utility. While at first glance Chen *et al.* offers some limited support for the PTO’s position, closer examination shows that the experimental design is flawed. In addition, any support to the PTO offered by Chen is offset by the authors’ citation of Celis *et al.* which supports Applicants’ position. Given the lack of support for the PTO’s position, Applicants submit that the PTO has not met its initial burden of overcoming the presumption that the asserted utility is credible and sufficient to satisfy the utility requirement. And even if the PTO has met that burden, the Applicants’ supporting rebuttal evidence is sufficient to establish



Appl. No. : 10/063,540  
Filed : May 2, 2002

that one of skill in the art would be more likely than not to believe that the claimed antibodies can be used as diagnostic tools for cancer, particularly esophageal and melanoma cancer.

### **Specific Utility**

#### *The Asserted Substantial Utilities are Specific to the Claimed Antibodies*

Applicants next address the PTO's assertion that the asserted utilities are not specific to the claimed antibodies related to PRO1277. Applicants respectfully disagree.

Specific Utility is defined as utility which is "specific to the subject matter claimed," in contrast to "a general utility that would be applicable to the broad class of the invention." M.P.E.P. § 2107.01 I. Applicants submit that the evidence of differential expression of the PRO1277 gene and polypeptide in certain types of tumor cells, along with the declarations and references discussed above, provide a specific utility for the claimed antibodies.

As discussed above, there are significant data which show that the gene for the PRO1277 polypeptide is expressed at least two-fold higher in normal esophageal and skin tissue compared to esophageal and melanoma tumor, respectively. These data are strong evidence that the PRO1277 gene and polypeptide are associated with esophageal and melanoma tumors. Thus, contrary to the assertions of the PTO, Applicants submit that they have provided evidence associating the PRO1277 gene and polypeptide with a specific disease. The asserted utility for antibodies to the PRO1277 polypeptide as a diagnostic tool for cancer, particularly esophageal and melanoma tumor, is a specific utility – it is not a general utility that would apply to the broad class of antibodies.

### **Conclusion**

The PTO has asserted three arguments to support its conclusion that based on the cited literature, one of skill in the art would not assume that higher expression of mRNA would correlate with increased polypeptide levels: (1) the PTO has challenged the reliability of the evidence reported in Example 18; (2) the PTO cites Pennica *et al.* to support its position that gene amplification is not necessarily correlated to gene expression; and (3) the PTO cites Hu *et al.*, Haynes *et al.* and Chen *et al.* to support its assertion that the literature cautions against drawing conclusions based on small changes in transcript expression levels between normal and

**Appl. No.** : 10/063,540  
**Filed** : May 2, 2002

cancerous tissue, and that mRNA levels are not predictive of protein levels. The PTO states that further research needs to be done to determine if the increase or decrease in PRO1277 DNA supports a role for the peptide in cancerous tissue. Applicants have addressed each of these arguments in turn.

First, the Applicants provided a first Declaration of Chris Grimaldi stating that the gene expression data in Example 18 are real and significant. This declaration also indicates that given the relative difference of at least two-fold in expression levels, the disclosed nucleic acids and corresponding polypeptides and antibodies have utility as cancer diagnostic tools. Hu *et al.* does not support the PTO's position, and is not contrary to Applicants' asserted utility. Thus, the PTO has not offered any substantial reason or evidence to question the data in Example 18, or the first Grimaldi Declaration.

Second, Applicants have shown that the second Grimaldi Declaration and Polakis Declaration, the accompanying references, as well as the excerpts and references cited above, demonstrate that it is well-established in the art that a change in mRNA levels generally correlates to a corresponding change in protein levels. Haynes *et al.* does not address this issue, and is not contrary to Applicants' asserted utility. Chen offers only very limited support for the PTO's position, and is not contrary to Applicants' assertion that generally there is a correlation. Thus, the PTO has not offered any substantial reason or evidence to question Applicants' declarations and supporting references.

Third, Applicants have shown that it is not necessary to know what role PRO1277 plays in cancer to use it as a diagnostic tool. The PTO's own guidelines recognize this fact, and numerous patents have issued which claim differentially expressed polypeptides and antibodies to the same, or methods employing such antibodies.

Finally, the PTO asserts that there is no asserted specific utility. Applicants have pointed out that the substantial utilities described above are specific to the claimed antibodies because the PRO1277 gene and polypeptide are differentially expressed in certain cancer cells compared to the corresponding normal cells. This is not a general utility that would apply to the broad class of antibodies.

Given the totality of the evidence provided, Applicants submit that they have established a substantial, specific, and credible utility for the claimed antibodies as diagnostic tools.

Appl. No. : 10/063,540  
Filed : May 2, 2002

According to the PTO Utility Examination Guidelines (2001), irrefutable proof of a claimed utility is not required. Rather, a specific, substantial, and credible utility requires only a “reasonable” confirmation of a real world context of use. Applicants remind the PTO that:

A small degree of utility is sufficient . . . The claimed invention must only be capable of performing some beneficial function . . . An invention does not lack utility merely because the particular embodiment disclosed in the patent lacks perfection or performs crudely . . . A commercially successful product is not required . . . Nor is it essential that the invention accomplish all its intended functions . . . or operate under all conditions . . . partial success being sufficient to demonstrate patentable utility . . . In short, **the defense of non-utility cannot be sustained without proof of total incapacity**. If an invention is only partially successful in achieving a useful result, a rejection of the claimed invention as a whole based on a lack of utility is not appropriate. M.P.E.P. at 2107.01 (underline emphasis in original, bold emphasis added, citations omitted).

Applicants submit that they have established that it is more likely than not that one of skill in the art would reasonably accept the utility for the claimed antibodies relating to PRO1277 set forth in the specification. In view of the above, Applicants respectfully request that the PTO reconsider and withdraw the utility rejection under 35 U.S.C. §101.

**Rejections under 35 U.S.C. § 112, first paragraph – Enablement**

The PTO also rejects Claims 1-5 under 35 U.S.C. § 112, first paragraph. Specifically, the PTO asserts that because the claimed invention is not supported by either a specific or substantial asserted utility or a well-established utility, one skilled in the art would not know how to use the claimed invention.

Applicants submit that in the discussion of the 35 U.S.C. § 101 rejection above, Applicants have established a substantial, specific, and credible utility for the claimed antibodies. To the extent that the enablement rejection is based on a lack of utility, Applicants respectfully request that the PTO reconsider and withdraw the enablement rejection under 35 U.S.C. §112.

Appl. No. : 10/063,540  
Filed : May 2, 2002

### CONCLUSION

In view of the above, Applicants respectfully maintain that claims are patentable and request that they be passed to issue. Applicants invite the Examiner to call the undersigned if any remaining issues may be resolved by telephone.

Please charge any additional fees, including any fees for additional extension of time, or credit overpayment to Deposit Account No. 11-1410.

Respectfully submitted,

KNOBBE, MARTENS, OLSON & BEAR, LLP

Dated: May 27, 2005

By: AnneMarie Kaiser  
AnneMarie Kaiser  
Registration No. 37,649  
Attorney of Record  
Customer No. 30,313  
(619) 235-8550

1674902  
042005